



# A Mutation-Independent CRISPR/Cas9-Based 'Knockout and Replace' Strategy to Treat Rhodopsin-Associated Autosomal Dominant Retinitis Pigmentosa (RHO-adRP)

*Zoe (Chi-Hsiu) Liu, Pavlina Wolf, Ruhong Dong, Yan Huang, Diana Tabbaa, Eugenio Marco, Brian Duke, Andrea Pinilla, Asha Pant, Racheal D'Souza, Judith Newmark, Georgia Giannoukos, Kate Zhang, Adrian Timmers, Mark S. Shearman and Mariacarmela Allocca*

***Presenter: Zoe (Chi-Hsiu) Liu***

May 4, 2022

# Disclosure

The speaker and the co-authors are employees and stockholders of Editas Medicine.

# Autosomal Dominant Retinitis Pigmentosa (adRP)

- An inherited autosomal dominant retinal disease leading to blindness in later life
- Symptoms:
  - Decreased night vision (nyctalopia)
  - Loss of peripheral vision (tunnel vision), and eventually significant decline in central vision
- No approved treatments





# Rhodopsin-Associated adRP (RHO-adRP)

## Rhodopsin (RHO)

- A light-sensitive receptor protein involved in visual phototransduction in rods
- Located in the outer segments of rods
- Approximately 30% (US and UK) of adRP caused by *RHO* dominant mutations
- Prevalence: 7,500 patients in US and 12,100 patients in EU and UK
- >150 mutations identified in the *RHO* gene cause RHO-adRP<sup>1</sup>
- Dominant mutations in the *RHO* gene are toxic for the rods: progressive loss of rods followed by loss of cones

## Photoreceptor structure



Valle D. et al. OMMBID 2014




# EDIT-103: Dual AAV-Based “Knockout and Replace” Therapeutic Strategy






- Agnostic to any *RHO* mutation – thus will knockout any dominant gain-of-function rhodopsin mutant
- **Step 1:** Both mutant and normal endogenous *RHO* will be knocked out in the treated area
- **Step 2:** Exogenous normal *RHO* (resistant to editing) will replace endogenous *RHO*
- **One-time subretinal administration aimed to restore/prevent vision loss**

## Details:

- gRNA is on the vector carrying *RHO-replace* thus assuring knockout takes place only in photoreceptor cells that express *RHO-replace*
- The *RHO* promoter for *Cas9* and *RHO-replace* restricts therapeutic activity to rod photoreceptors




# The RHO Promoter Restricts Gene Expression to Rod Photoreceptors in the Mouse Eyes



Scale bar = 20  $\mu$ m


# EDIT-103 is Highly Specific: No Detectable Off-Target Editing

**NO OFF-TARGET EDITING AT OFF-TARGET CANDIDATES**  
(rhAmpSeq verification in human retina explants transduced with EDIT-103)




# EDIT-103 in Humanized $mRho^{hRHO/+}$ Mice: Demonstrates Rapid and Stable Gene Editing




Dose escalation



Time course






**$hRHO$  mRNA replacement**



- >25% of rods are edited
- Editing rates increase in a dose-dependent manner
- Editing plateau at ~6 weeks post-injection at doses  $\geq 3E12$  vg/ml

AAV ratio: 1:1  
Volume: 1  $\mu$ l  
Geometric mean with  
95% confidence interval is presented

# EDIT-103 in Non-Human Primates (NHPs): Approximately 100% Editing in Transduced Photoreceptors



~100% editing (within the transduced area) in NHP





# EDIT-103 in NHPs: Nearly Complete Knockdown of the Endogenous RHO and Over 30% RHO Protein Replacement



Endogenous NHP RHO

Human RHO Replacement

*RHO* mRNA



*RHO* Protein




- 80% and 90% knockdown of endogenous NHP *RHO* mRNA levels were achieved at doses of 3E12 and 6E12 vg/ml, respectively. This resulted in 90% and 100% of *RHO* protein knockdown.
- RHO* replacement mRNA levels increased with dose and resulted in >30% of *RHO* protein levels.

Volume: 100  $\mu$ l  
AAV ratio: 1:1  
Time point: 13 weeks  
Mean ( $\pm$ SD) is presented  
\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$



AAV: adeno-associated virus; coRHO: codon-optimized rhodopsin; KO: knockout; mRNA: messenger ribonucleic acid; NHP: non-human primate; RHO: rhodopsin; SD: standard deviation.

# RHO Protein Expression and Retina Morphology was Preserved in EDIT-103-Treated NHP Retinas Compared with KO-Treated Retinas




# Retina Function Preserved in the EDIT-103-Treated NHP Eyes Compared to the KO Only Treated Eyes

Retinal layers



AAV ratio: 1:1; Injection volume: 100  $\mu$ L; Time point: 13 weeks; Mean ( $\pm$ SD) is presented; \* $p$ <0.05, \*\*\* $p$ <0.001

Dark-adapted 3.0 ERG  
(combined rod-cone response)



- KO of endogenous *RHO* significantly reduced *a*- and *b*-wave amplitudes
- EDIT-103 dosing preserved *a*- and *b*-wave amplitudes

# Summary



EDIT-103 is a **one-time, high efficacy, mutation-agnostic gene medicine** to permanently suppress the toxic gain-of-function associated with RHO-adRP



**Ex vivo:** EDIT-103 shows high specificity in human retinal explants



**In vivo:**

- *mRho*<sup>hRHO/+</sup> mouse:
  - EDIT-103 achieved **rapid** and **stable** gene editing:
    - Editing plateau at 6 weeks and is sustained until end of study (13 weeks)
    - > 25% gene editing at doses  $\geq 3E12$  vg/ml
- NHP:
  - EDIT-103 achieved nearly **100%** editing
  - **>30%** RHO replacement protein levels
  - Morphological and functional photoreceptor preservation

# Acknowledgements

Mariacarmela Allocca

Ruhong Dong

Racheal D'Souza

Brian Duke

Georgia Giannoukos

Yan Huang

Eugenio Marco

Judith Newmark

Asha Pant

Andrea Pinilla

Mark S. Shearman

Diana Tabbaa

Adrian Timmers

Pavlina Wolf

Kate Zhang

\* Alphabetical order according to last names



Thank you

